

MIDAZOLAM (Versed)

ACTION: Hypnotic, Sedative, Anti-Seizure

- Midazolam is a potent, short-acting benzodiazepine with hypnotic and amnestic effects. It has no effect on pain.

INDICATIONS:

- Premedication before cardioversion, external pacing and other painful procedures.
- Seizures (status epilepticus).
- Agitated patient who may be a danger to self or others.

CONTRAINDICATIONS:

- Hypersensitivity
- Shock, with depressed vital signs
- Narrow-angle glaucoma
- Alcoholic coma

POTENTIAL SIDE EFFECTS:

- Laryngospasm
- AMS
- Bronchospasm
- Bradycardia
- Dyspnea
- Tachycardia
- Respiratory depression and arrest
- PVC's
- Drowsiness
- Nausea
- Amnesia
- Vomiting

ADULT DOSE/ROUTE:

- ⇒ **Sedation/Agitation:** 2 - 5mg IM x1 or 1 – 2mg slow push IV/IO. May repeat IV/IO dose in 5 minutes for continued agitation. Maximum total dose 5mg IV/IO.
- ⇒ **Seizure:** 10 mg IM x1 or intranasally (5mg each nostril) or 5mg slow push IV/IO. May repeat IV/IO dose in 5 minutes. Maximum total dose 10mg IV/IO.

PEDIATRIC DOSE/ROUTE:

- ⇒ **Sedation/Agitation:** 0.2 mg/kg IV/IM/IO or 0.4 mg/kg intranasally to a maximum single dose of 5mg. May repeat the initial weight-based dose x1. Maximum total dose should not exceed 10 mg.
- ⇒ **Seizure:** 0.2 mg/kg IV/IM/IO or 0.4 mg/kg intranasally to a maximum single dose of 5mg. May repeat the initial weight-based dose x1. Maximum total dose should not exceed 10 mg.

NOTES:

- Contact Base Hospital if seizure activity continues after maximum dose is reached and additional doses are required.
- Do not use intranasal route in agitation because amount of absorption in an actively resisting, agitated patients is unknown.
- ~~Midazolam is more potent than diazepam.~~
- Effects of midazolam may be potentiated if administered with morphine opiate pain medications. Contact Base Hospital if considering administering both medications.

- Always be prepared to protect airway and ventilate patients who are given benzodiazepines. Continuous monitoring of vital signs before and after administration is required.

DRAFT